



# Amino acid $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ from sclerotized beaks: a new tool to investigate the foraging ecology of cephalopods, including giant and colossal squids

Yves Cherel<sup>1,\*</sup>, Paco Bustamante<sup>2</sup>, Pierre Richard<sup>2</sup>

<sup>1</sup>Centre d'Etudes Biologiques de Chizé (CEBC), UMR 7372 du CNRS-La Rochelle Université, 79360 Villiers-en-Bois, France

<sup>2</sup>Littoral, Environnement et Sociétés (LIENSs), UMR 7266 du CNRS- La Rochelle Université, 2 rue Olympe de Gouges, 17000 La Rochelle, France

**ABSTRACT:** Combining the use of predators as biological samplers together with measurements of the stable isotopic ratios ( $\delta^{13}\text{C}_{\text{Bulk}}$  and  $\delta^{15}\text{N}_{\text{Bulk}}$ ) of their sclerotized beaks help investigate foraging ecology of poorly known oceanic cephalopods. However, high chitin content (an amino-sugar macromolecule) lowers beak  $\delta^{15}\text{N}_{\text{Bulk}}$  values, thus precluding direct isotopic comparison with other tissues and organisms. To overcome the chitin effect, compound-specific isotopic analysis of amino acids (CSIA-AA) was performed on squid beaks. The method was applied on beaks and muscle, and the resulting  $\delta^{13}\text{C}_{\text{AA}}$  and  $\delta^{15}\text{N}_{\text{AA}}$  values compared between tissues. The usefulness of CSIA was tested by defining the habitat and trophic position ( $\text{TP}_{\text{CSIA}}$ ) of squids using their  $\delta^{13}\text{C}_{\text{AA}}$  and  $\delta^{15}\text{N}_{\text{AA}}$  values. Beak  $\delta^{13}\text{C}_{\text{AA}}$  values were reliably measured on 12 AA that included 5 essential and 7 non-essential AA, and  $\delta^{15}\text{N}_{\text{AA}}$  values were quantified on at least 7 AA that included 2 source and 4 trophic AA. Importantly,  $\delta^{13}\text{C}_{\text{AA}}$  and  $\delta^{15}\text{N}_{\text{AA}}$  varied little between muscle and lower and upper beaks, and  $\text{TP}_{\text{CSIA}}$  estimates were identical regardless of the tissue considered. Tissue  $\delta^{13}\text{C}_{\text{AA}}$  values of both essential and non-essential AA reflected the latitudinal baseline  $\delta^{13}\text{C}$  gradient that occurs in the Southern Indian Ocean, while beak  $\delta^{15}\text{N}_{\text{AA}}$  from source and trophic AA allowed the disentangling of the baseline effect from the trophic effect, and thus better calculations of squid TP estimates than from  $\delta^{15}\text{N}_{\text{Bulk}}$  values. Beak  $\delta^{13}\text{C}_{\text{AA}}$  and  $\delta^{15}\text{N}_{\text{AA}}$  defined isotopic niches of colossal and giant squids, the 2 largest living invertebrates. In subantarctic waters, they segregate by having species-specific foraging habitats (using  $\delta^{13}\text{C}_{\text{Gly}}$  or  $\delta^{15}\text{N}_{\text{Phe}}$ ) and  $\text{TP}_{\text{CSIA}}$  (using  $\delta^{15}\text{N}_{\text{Glx}}$  and  $\delta^{15}\text{N}_{\text{Phe}}$ ).  $\text{TP}_{\text{CSIA}}$  is higher in colossal (4.7) than giant (4.3) squids, and both values compare well with those of myctophid-eaters, suggesting very large squids prey primarily upon small zooplanktivorous fishes. As expected, CSIA-AA overcomes the chitin effect on beaks and it is a powerful tool to investigate trophic interactions of cephalopods. The method has a great potential with arthropods, because chitin is a main component of their exoskeleton but the deleterious effect of chitin is overlooked in isotopic studies focusing on crustaceans and insects.

**KEY WORDS:** Arthropod · Carbon · Chitin · CSIA · Habitat · Nitrogen · Stable isotope · Trophic position

## 1. INTRODUCTION

Cephalopods play a major role in marine trophic webs, as underlined by their global biomass and annual consumption of resources (Clarke 1996, Coll et al. 2013). Determining and quantifying their trophic

relationships is therefore key to understanding the structure and functioning of marine ecosystems. The role of cephalopods as prey is demonstrated by their importance in the diet of predators, but knowledge of their food is limited by lack of data (Clarke 1996). A new approach to investigate cephalopod feeding

\*Corresponding author: cherel@cebc.cnrs.fr

ecology was developed by combining the use of their predators as biological samplers together with measurements of the stable isotopic values of their beaks (Cherel & Hobson 2005, Cherel et al. 2009b). Predators as samplers have 2 main advantages: (1) they catch larger specimens and a greater diversity of cephalopods than fisheries sampling gear (Rodhouse 1990), and (2) they accumulate hundreds to thousands of beaks in their stomachs, because beaks are hard sclerotized structures that resist digestion (Clarke 1980, Cherel et al. 2017).

The basic concept of the stable isotope method is that an animal's isotopic composition is directly influenced by the food it assimilates. The 2 main elements used in isotopic ecology are carbon and nitrogen, whose isotopic ratios are measured on bulk tissue (mainly muscle) that contains primarily proteins. Consumer proteins are enriched in  $^{15}\text{N}$  relative to dietary proteins, and consequently  $\delta^{15}\text{N}_{\text{Bulk}}$  measurements serve as indicators of a consumer's diet and trophic position ( $\text{TP}_{\text{Bulk}}$ ) (Vanderklift & Ponsard 2003). By contrast,  $\delta^{13}\text{C}_{\text{Bulk}}$  varies little along the food web and is mainly used to determine primary sources in a trophic network (Kelly 2000). Measuring  $\delta^{13}\text{C}_{\text{Bulk}}$  and  $\delta^{15}\text{N}_{\text{Bulk}}$  on beaks gave new insights in cephalopod biology, such as latitudinal feeding habitats, migration patterns,  $\text{TP}_{\text{Bulk}}$ , trophic structure of the communities, and ontogenic dietary changes (Cherel & Hobson 2005, Cherel et al. 2009a,b, Navarro et al. 2013, Golikov et al. 2018). However, while  $\delta^{13}\text{C}_{\text{Bulk}}$  values of beaks and soft tissues are similar, beaks have consistently lower  $\delta^{15}\text{N}_{\text{Bulk}}$  values (Hobson & Cherel 2006, Ruiz-Cooley et al. 2006, Cherel et al. 2009a). This major limitation precludes comparing raw (uncorrected) beak  $\delta^{15}\text{N}_{\text{Bulk}}$  values with those of other tissues and organisms to trace trophic pathways in marine ecosystems (Hobson & Cherel 2006, Cherel et al. 2009a).

Beaks and soft tissues have different biochemical compositions. Beaks contain chitin (Hunt & Nixon 1981, Rubin et al. 2010), a modified polysaccharide that is impoverished in  $^{15}\text{N}$  compared to consumer diet (Schimmelmann 2011). The presence of chitin explains why beaks have lower  $\delta^{15}\text{N}_{\text{Bulk}}$  values than soft tissues. Moreover, the ratio of chitin to protein varies within beaks and between beaks, because the undarkened, darkening and darkened parts of beaks contain decreasing relative amounts of chitin over protein (Rubin et al. 2010). Hence, overcoming the chitin effect on beaks is of primary importance, and 4 different approaches can be used theoretically (Xavier et al. 2015). (1) Quantification of isotopic correction factors allows the comparison of  $\delta^{15}\text{N}_{\text{Bulk}}$  be-

tween beaks and soft tissues, with the drawback that corrected values are estimates (Hobson & Cherel 2006, Cherel et al. 2009a). (2) The same limitation applies to chitin normalization models using C:N mass ratios as a proxy for chitin content, following similar methods applied to correct for variable lipid content in bulk  $\delta^{13}\text{C}$  analyses (as lipids, chitin has a higher C:N value than proteins; Webb et al. 1998). (3) Measurements of  $\delta^{15}\text{N}$  on purified proteins is not feasible because classical extraction protocols for soft tissues are not effective for beaks (the majority of proteins remain insoluble even under the most aggressive extraction procedures; Rubin et al. 2010). (4) Measurements of  $\delta^{15}\text{N}$  on amino acids (AA) from protein is a promising tool, but no studies used the compound-specific isotopic analysis of amino acids (CSIA-AA) (McMahon & McCarthy 2016) on cephalopod beaks.

CSIA-AA has emerged in the last decade as a powerful approach for tracing the origins and fate of carbon and nitrogen in ecological and biogeochemical studies (McMahon et al. 2013, Ohkouchi et al. 2017). The method has a broad range of applications, including the identification of baseline isoscapes, the assessment of the source and transformation of detrital organic matter, and tracing of animal migration. While comparatively few investigations are based on  $\delta^{13}\text{C}_{\text{AA}}$  measurements,  $\delta^{15}\text{N}_{\text{AA}}$  values are increasingly used to calculate accurate trophic position estimates ( $\text{TP}_{\text{CSIA}}$ ) of a broad range of terrestrial and aquatic consumers (Chikaraishi et al. 2014). To our knowledge, no study has measured  $\delta^{13}\text{C}_{\text{AA}}$  in cephalopods, and only limited information is available on their  $\delta^{15}\text{N}_{\text{AA}}$  values. Indeed, a preliminary study shows incidentally a chromatogram from nitrogen analysis of a single squid beak (Walsh et al. 2014), but most previous  $\delta^{15}\text{N}_{\text{AA}}$  measurements have been restricted to the mantle of a few specimens of ommastrephid squids (Ruiz-Cooley et al. 2013, Madigan et al. 2016, Hetherington et al. 2017).

The main goal of the present study was to use CSIA-AA on cephalopod beaks primarily to bypass the chitin effect. We focused on 3 points: (1) we validated how many and which AA can be isolated from beaks to reliably measure their  $\delta^{13}\text{C}_{\text{AA}}$  and  $\delta^{15}\text{N}_{\text{AA}}$  values. (2) Using the same specimens, we compared  $\delta^{13}\text{C}_{\text{AA}}$  and  $\delta^{15}\text{N}_{\text{AA}}$  values of lower and upper beaks with those of muscle, the canonical tissue for isotopic investigations. We hypothesized that AA isotopic values are identical in muscle (a metabolically active tissue) and in the more recently built parts of beaks (a metabolically inactive tissue). (3) We investigated the biological usefulness of CSIA-AA on beaks by testing

(i) if beak  $\delta^{13}\text{C}_{\text{AA}}$  reflects the latitudinal baseline  $\delta^{13}\text{C}$  gradient occurring in the Southern Indian Ocean; the gradient allows defining the latitudinal habitat of consumers using either  $\delta^{13}\text{C}_{\text{Bulk}}$  (Jaeger et al. 2010) or  $\delta^{13}\text{C}_{\text{AA}}$  (Lorrain et al. 2009); and (ii) if beak  $\delta^{13}\text{C}_{\text{AA}}$  and  $\delta^{15}\text{N}_{\text{AA}}$  help to define the isotopic niche (feeding habitat and  $\text{TP}_{\text{CSIA}}$ ) and thus the mechanisms allowing co-existence of 4 sympatric cephalopods that include the 2 largest living invertebrates, the colossal squid *Mesonychoteuthis hamiltoni* and giant squid *Architeuthis dux*.

## 2. MATERIALS AND METHODS

### 2.1. Study sites, dietary sampling and analysis

Fieldwork was carried out in the Southern Indian Ocean by fishery observers during commercial cruises. The fishery targeted Patagonian toothfish *Dissostichus eleginoides*, with southern sleeper sharks *Somniosus antarcticus* being occasionally by-caught. Cephalopod items were sorted from fish stomachs and kept in 70% ethanol until analysis. Squid lower beaks were identified from their morphological features by comparison with material held in our own collection and by reference to the available literature (Xavier & Cherel 2009). Lower rostral length of beaks were measured to 0.1 mm with a vernier caliper and allometric equations were used to estimate dorsal mantle length (ML) of squids (Table 1).

Two sets of samples were analyzed. (1) Ten whole buccal masses of the giant warty squid *Kondakovia longimana* were collected from toothfish stomachs in

Kerguelen ( $n = 7$ ) and Crozet waters ( $n = 3$ ) in 2014–2015. These were dissected to sort lower and upper beaks from buccal masses. Three tissues were sampled for isotopic analysis: muscle tissue, wings of lower beaks, and small pieces of lateral walls of upper beaks. (2) Accumulated beaks from stomachs of toothfish and sharks caught in Kerguelen waters in 1997–2001 (Cherel & Duhamel 2004, Cherel et al. 2004) were used to compare the isotopic values of *M. hamiltoni*, *A. dux*, *K. longimana*, and Dana octopus squid *Taningia danae*. Wings or free lateral corners of lateral walls from lower beaks were cut with scissors. Importantly, all the different sampled parts of lower and upper beaks referred to newly built material, to minimize potential trophic ontogenetic changes (Cherel & Hobson 2005, Queiros et al. 2018) and different tissue-related time integration periods between beaks and the metabolically active muscle tissue.

### 2.2. Stable isotope analysis

Beaks were cleaned before analysis to remove any remains of soft tissue and mucus. Samples were freeze-dried and ground to a fine powder. Bulk and AA  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values were determined on the same samples.

**2.2.1. Bulk isotopic measurements.** Lipids of muscle tissue were removed using cyclohexane. Subsamples of the homogenates of beaks and of lipid-extracted muscle tissue were weighed with a microbalance and packed in tin cups. An elemental analyser (Thermo Scientific Flash 2000) was coupled to a continuous

Table 1. Measured lower rostral length (LRL) and estimated mantle length (ML) of oceanic squids from Kerguelen waters. Predator: species from which the beaks were extracted. ML was calculated using species-specific allometric equations from the corresponding references. Values are means  $\pm$  SD (parentheses: range). Kruskal-Wallis  $H$ -tests and Conover-Inman tests for pairwise comparisons were performed to compare values amongst the 4 species (values in the same column with differing superscript letters are statistically different). Significant differences ( $p < 0.05$ ) are highlighted in **bold**

Squid species	Predator	n	LRL (mm)	ML (cm)	Reference
<i>Architeuthis dux</i> (giant squid)	<i>Somniosus antarcticus</i> (southern sleeper shark)	6	15.3 $\pm$ 3.0 <sup>a</sup> (11.3–18.0)	141 $\pm$ 72 <sup>a</sup> (54–215)	Roeleveld (2000)
<i>Kondakovia longimana</i> (giant warty squid)	<i>Dissostichus eleginoides</i> (Patagonian toothfish)	15	13.9 $\pm$ 1.7 <sup>a</sup> (10.5–17.8)	50 $\pm$ 6 <sup>b</sup> (37–64)	Adams & Klages (1987)
<i>Mesonychoteuthis hamiltoni</i> (colossal squid)	<i>Somniosus antarcticus</i> (southern sleeper shark)	10	29.2 $\pm$ 5.8 <sup>b</sup> (23.2–39.0)	178 $\pm$ 35 <sup>a</sup> (141–239)	Clarke (1986)
<i>Taningia danae</i> (Dana octopus squid)	<i>Somniosus antarcticus</i> (southern sleeper shark)	10	18.7 $\pm$ 1.8 <sup>c</sup> (15.5–21.0)	85 $\pm$ 13 <sup>c</sup> (61–102)	Clarke (1986)
Kruskal-Wallis $H$ -test	$H$		<b>31.4</b>	<b>31.6</b>	
	$p$		<b>&lt;0.0001</b>	<b>&lt;0.0001</b>	

flow mass spectrometer (Thermo Scientific Delta V Plus) to measure carbon and nitrogen contents, and  $\delta^{13}\text{C}_{\text{Bulk}}$  and  $\delta^{15}\text{N}_{\text{Bulk}}$  values, respectively. Stable isotope ratios are expressed using standard  $\delta$  notation relative to carbonate Vienna PeeDee Belemnite and atmospheric nitrogen. Two internal standards of caffeine (USGS 61 and USGS 62) were used for drift assessment and data normalization. Observed analytical errors on internal standards were  $<0.10\%$  for both  $\delta^{13}\text{C}_{\text{Bulk}}$  and  $\delta^{15}\text{N}_{\text{Bulk}}$ .

**2.2.2. CSIA-AA.** Beak and non-lipid extracted muscle samples (1–2 mg) were hydrolyzed under nitrogen (0.5 ml 6 M HCl, 110°C, 20 h). Norleucine (20  $\mu\text{l}$ , 25 mM) was added to each sample as an internal standard prior to hydrolysis. The resultant AA were purified and derivatized to N-acetyl-isopropyl esters (Styring et al. 2012). Caffeine (IAEA-600) was added as an internal standard to the derivatized AA before dilution in ethyl acetate for carbon and nitrogen analyses by continuous-flow gas chromatography-combustion-isotope ratio mass spectrometry (GC-C-IRMS).  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values were measured using a Thermo Trace GC Ultra gas chromatograph coupled to a Delta V Plus isotope-ratio mass spectrometer via a GC IsoLink II interface (Thermo Scientific). The combustion/reduction reactor was maintained at 1000°C, and a liquid nitrogen cold trap was used after the reactor to remove  $\text{CO}_2$  during nitrogen analyses. AA were separated on a VF-35MS column (30 m, 0.32 mm ID, 1  $\mu\text{m}$  film thickness; Agilent Technologies). Analyses were done with a splitless injection at 270°C, and a helium flow set at 1.4 ml  $\text{min}^{-1}$ . Samples were analyzed either in duplicate or triplicate. A mixture of 16 AAs and Norleucine, thoroughly calibrated by EA-IRMS and derivatized along with the samples, was injected after every 4 samples to evaluate drift and accuracy. Raw data were corrected (Docherty et al. 2001) and normalized using internal standard values (norleucine:  $\delta^{13}\text{C}$ :  $-28.77 \pm 0.05\%$ ,  $\delta^{15}\text{N}$ :  $19.19 \pm 0.08\%$ ; caffeine:  $\delta^{13}\text{C}$ :  $-27.77 \pm 0.04\%$ ,  $\delta^{15}\text{N}$ :  $1.00 \pm 0.20\%$ ). Depending on AA, measurement precision for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  of the standard mixture ranged from 0.2 to 1.0‰ (mean 0.4‰), and from 0.2 to 1.2‰ (mean 0.5‰), respectively.

### 2.3. General comments about AA

Twenty AAs form the essential building blocks of proteins. Acid hydrolysis destroys tryptophan and precludes determining cysteine directly (Fountoulakis & Lahm 1998). It also converts asparagine (Asn) and glutamine (Gln) into aspartic acid (Asp) and glu-

tamic acid (Glu), respectively, resulting in the measurements of combined Asn + Asp (Asx) and Gln + Glu (Glx). Since arginine is not derivatized, the analytical procedure allows the quantification of  $\delta^{13}\text{C}_{\text{AA}}$  and  $\delta^{15}\text{N}_{\text{AA}}$  for 15 standard AAs at best, because some AAs (Met, Ser, Thr, Tyr; see abbreviations in Table 2) are partially destroyed by hydrolysis (Fountoulakis & Lahm 1998). In animals, standard AAs are classified into 2 categories (essential or non-essential) with regard to carbon metabolism. Eight of the 15 measured AAs (His, Ile, Leu, Lys, Met, Phe, Thr and Val) are essential AAs, with the remaining 7 AAs being non-essential (Ala, Asx, Glx, Gly, Pro, Ser and Tyr) (Lehninger 1982). In terms of  $\delta^{15}\text{N}$ , the 15 AAs group into 5 source AAs (His, Lys, Phe, Met and Tyr) and 7 trophic AAs (Ala, Asx, Glx, Ile, Leu, Pro and Val). Trophic AAs undergo significant  $^{15}\text{N}$  enrichment between food and consumers, while source AAs do not, thus reflecting  $\delta^{15}\text{N}$  baseline. Gly and Ser are 2 challenging AAs to classify into the source and trophic framework, and were clustered into a source/trophic group. Finally, Thr is considered as a metabolic AA because it shows  $^{15}\text{N}$  depletion relative to dietary Thr (McMahon & McCarthy 2016).

### 2.4. Trophic position and data analyses

Lower beak  $\delta^{15}\text{N}_{\text{AA}}$  values of the 2 canonical source AA Phe and trophic AA Glx (McMahon & McCarthy 2016) were used to estimate  $\text{TP}_{\text{CSIA}}$  and calculate the relative trophic position (RTP) of squids. The first formulation is based on equations from Chikaraishi et al. (2010) and McMahon & McCarthy (2016), as:

$$\text{TP}_{\text{Glx-Phe}} = [(\delta^{15}\text{N}_{\text{Glx}} - \delta^{15}\text{N}_{\text{Phe}} - \text{TDF}_1 - \beta) / \text{TDF}_2] + 2$$

where  $\text{TDF}_1$  represents the trophic discrimination factor ( $\text{TDF}_{\text{Glx-Phe}}$ ) between food and consumers typical of lower trophic-level organisms (7.6‰; Chikaraishi et al. 2010),  $\beta$  is the difference in  $\delta^{15}\text{N}$  between Glx and Phe in primary producers at the base of the food web (2.9‰; Nielsen et al. 2015) and  $\text{TDF}_2$  reflects  $\text{TDF}_{\text{Glx-Phe}}$  for cephalopods (5.0‰; McMahon & McCarthy 2016). The second formulation is a proxy for TP: since  $\text{RTP} = \delta^{15}\text{N}_{\text{Glx}} - \delta^{15}\text{N}_{\text{Phe}}$  of consumers, it is expressed in ‰, thus differing from TP, which is a rational number from 1 to up to 6 (with no unit). RTP calculation requires no *a priori* assumptions about  $\beta$  and TDF values used to estimate  $\text{TP}_{\text{Glx-Phe}}$  other than the assumption that these values remain constant among the samples. RTP essentially removes the isotopic effect of food web baseline, focusing on relative differences in food web position (Choy et al. 2015).

Estimated TP of squids was also calculated using  $\delta^{15}\text{N}_{\text{Bulk}}$  values of their lower beaks (modified from Cherel et al. 2008), as  $\text{TP}_{\text{Bulk}} = [(\delta^{15}\text{N}_{\text{Bulk}} + 0.10) / \text{TDF}] + 2$ , where 0.10 is the difference between 3.46‰ (the isotopic factor to correct the chitin effect between wings of lower beaks and muscle tissue; Cherel et al. 2009a) and 3.36‰ (the average  $\delta^{15}\text{N}_{\text{Bulk}}$  value of the herbivorous salp *Salpa thompsoni* in Kerguelen waters with an assumed TP of 2.0; Cherel et al. 2008, 2010), and TDF is the  $\delta^{15}\text{N}$  difference between muscle of cephalopods and their food (3.3‰; Hobson & Cherel 2006).

Estimating TP is challenging, with every method showing limitations. Three major issues are (1) quantification of TDF for  $\text{TP}_{\text{Bulk}}$  and  $\text{TP}_{\text{CSIA}}$ , (2)  $\beta$  for  $\text{TP}_{\text{CSIA}}$ , and (3) baseline value for  $\text{TP}_{\text{Bulk}}$ . In aquatic ecosystems, particulate organic matter (POM) is often used as a proxy for phytoplankton and as a food-web baseline for  $\text{TP}_{\text{Bulk}}$  calculations (Post 2002). However, the use of POM is not ideal, because (1) it represents an unknown mixture of phytoplankton together with detritus, bacteria and microzooplankton, and (2) its turnover is high, thus promoting large  $\delta^{15}\text{N}_{\text{Bulk}}$  variations at small temporal scales that are buffered in higher TP organisms (Pakhomov et al. 2019). An alternative to POM is to consider longer-lived primary consumers (herbivorous copepods or pelagic

tunicates) that are assumed to be representative of  $\text{TP}_{\text{Bulk}} = 2.0$ . However, crustacean exoskeleton contains chitin that is likely to lower their  $\delta^{15}\text{N}_{\text{Bulk}}$  values (see Section 4.3). Hence, we used salps as control organisms in the present investigation, even though salps themselves are not always appropriate due to their selective feeding habits (Kruse et al. 2015, Pakhomov et al. 2019).

Data were statistically analyzed using SYSTAT 13. Values are means  $\pm$  SD.

### 3. RESULTS

#### 3.1. Comparing $\delta^{13}\text{C}_{\text{AA}}$ and $\delta^{15}\text{N}_{\text{AA}}$ in muscle and beaks

**3.1.1. Bulk  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values.** Bulk  $\delta^{13}\text{C}$  values of muscle, lower beak and upper beak of warty squids were not statistically different (Table 2). As expected, a chitin effect was found in  $\delta^{15}\text{N}_{\text{Bulk}}$  values, which were lower in beaks than in muscle tissue (Table 3).

**3.1.2. Amino acid  $\delta^{13}\text{C}$  values.** The analytical method quantified  $\delta^{13}\text{C}_{\text{AA}}$  of 16 AAs in muscle and 12 AAs in beaks. Too low amounts of 4 AA (Hyp, Ile, Lys and Met) precluded reliable isotopic measurements

Table 2. Bulk and individual amino acid  $\delta^{13}\text{C}$  values (‰) of muscle, lower beak and upper beak from buccal masses of giant warty squids. Values are means  $\pm$  SD. Kruskal-Wallis  $H$ -tests and Conover-Inman tests for pairwise comparisons were performed to compare  $\delta^{13}\text{C}$  values from the 3 tissues (values in the same row with differing superscript letters are statistically different). Significant differences ( $p < 0.05$ ) are highlighted in **bold**. (–) No data (see Section 3.1.2)

Amino acid	Abbreviation	Muscle	Lower beak	Upper beak	Kruskal-Wallis $H$	$H$ -test $p$
Bulk		$-22.9 \pm 1.6^a$	$-23.0 \pm 1.9^a$	$-22.7 \pm 1.9^a$	0.32	0.852
<b>Essential</b>						
Histidine	His	$-12.0 \pm 2.6^a$	$-11.1 \pm 2.4^a$	$-11.7 \pm 3.7^a$	0.10	0.952
Isoleucine	Ile	$-24.3 \pm 2.6$	–	–	–	–
Leucine	Leu	$-32.1 \pm 3.4^a$	$-32.0 \pm 3.7^a$	$-32.0 \pm 3.7^a$	0.07	0.967
Lysine	Lys	$-21.9 \pm 2.1$	–	–	–	–
Methionine	Met	$-28.1 \pm 2.5$	–	–	–	–
Phenylalanine	Phe	$-30.1 \pm 2.6^a$	$-28.8 \pm 2.7^a$	$-28.7 \pm 2.7^a$	1.68	0.433
Threonine	Thr	$-13.6 \pm 1.6^a$	$-11.7 \pm 1.6^b$	$-11.6 \pm 2.1^b$	<b>6.13</b>	<b>0.047</b>
Valine	Val	$-23.9 \pm 2.3^a$	$-24.0 \pm 2.7^a$	$-24.7 \pm 3.1^a$	0.73	0.693
<b>Non essential</b>						
Alanine	Ala	$-22.5 \pm 3.4^a$	$-22.2 \pm 3.6^a$	$-22.2 \pm 3.6^a$	0.07	0.967
Aspartic acid	Asx=Asn+Asp	$-19.4 \pm 2.9^a$	$-18.4 \pm 3.5^a$	$-18.7 \pm 3.6^a$	0.88	0.644
Glutamic acid	Glx=Gln+Glu	$-19.6 \pm 2.6^a$	$-18.1 \pm 2.8^a$	$-18.2 \pm 2.9^a$	1.79	0.409
Glycine	Gly	$-4.6 \pm 4.2^a$	$-5.8 \pm 4.2^a$	$-5.6 \pm 4.0^a$	0.87	0.647
Hydroxyproline	Hyp	$-20.2 \pm 2.3$	–	–	–	–
Proline	Pro	$-21.9 \pm 2.4^a$	$-21.3 \pm 3.3^a$	$-21.4 \pm 3.2^a$	0.35	0.839
Serine	Ser	$-5.1 \pm 3.5^a$	$-6.2 \pm 3.7^a$	$-5.4 \pm 4.9^a$	0.47	0.791
Tyrosine	Tyr	$-28.5 \pm 2.3^a$	$-25.6 \pm 3.0^a$	$-25.8 \pm 3.4^a$	4.07	0.131

Table 3. Bulk and individual amino acid  $\delta^{15}\text{N}$  values (‰) of muscle, lower beak and upper beak from buccal masses of giant warty squids. Values are means  $\pm$  SD. Kruskal-Wallis  $H$ -tests and Conover-Inman tests for pairwise comparisons were performed to compare  $\delta^{15}\text{N}$  values from the 3 tissues (values in the same row with differing superscript letters are statistically different). Significant differences ( $p < 0.05$ ) are highlighted in **bold**. TP: trophic position; RTP: relative trophic position (‰); (–) no data (see Section 3.1.3)

Amino acid	Group	Muscle	Lower beak	Upper beak	Kruskal-Wallis $H$ -test	$p$
Bulk		9.0 $\pm$ 0.4 <sup>a</sup>	6.5 $\pm$ 0.7 <sup>b</sup>	4.6 $\pm$ 1.0 <sup>c</sup>	<b>24.5</b>	<b>&lt;0.0001</b>
Alanine	Trophic	21.7 $\pm$ 2.0 <sup>a</sup>	21.8 $\pm$ 1.6 <sup>a</sup>	22.3 $\pm$ 1.3 <sup>a</sup>	0.11	0.945
Aspartic acid	Trophic	15.5 $\pm$ 1.9 <sup>a</sup>	17.9 $\pm$ 1.9 <sup>b</sup>	18.1 $\pm$ 2.0 <sup>b</sup>	<b>7.44</b>	<b>0.024</b>
Glutamic acid	Trophic	21.6 $\pm$ 0.7 <sup>a</sup>	21.7 $\pm$ 1.5 <sup>a</sup>	21.8 $\pm$ 1.6 <sup>a</sup>	0.56	0.756
Glycine	Source/trophic	–1.0 $\pm$ 1.1 <sup>a</sup>	–3.5 $\pm$ 2.0 <sup>b</sup>	–3.4 $\pm$ 2.3 <sup>b</sup>	<b>9.25</b>	<b>0.010</b>
Histidine	Source	6.2 $\pm$ 2.2 <sup>a</sup>	6.7 $\pm$ 1.9 <sup>a</sup>	6.2 $\pm$ 1.7 <sup>a</sup>	0.49	0.784
Isoleucine	Trophic	23.0 $\pm$ 1.2	–	–	–	–
Leucine	Trophic	22.4 $\pm$ 2.3 <sup>a</sup>	23.2 $\pm$ 2.3 <sup>a</sup>	23.5 $\pm$ 2.5 <sup>a</sup>	1.24	0.537
Lysine	Source	2.6 $\pm$ 0.6	–	–	–	–
Phenylalanine	Source	–1.0 $\pm$ 1.6 <sup>a</sup>	–0.5 $\pm$ 0.8 <sup>a</sup>	–1.0 $\pm$ 0.7 <sup>a</sup>	2.51	0.285
Proline	Trophic	22.8 $\pm$ 1.2 <sup>a</sup>	19.3 $\pm$ 3.2 <sup>b</sup>	20.7 $\pm$ 2.3 <sup>b</sup>	<b>8.47</b>	<b>0.014</b>
Serine	Source/trophic	5.4 $\pm$ 1.7 <sup>a</sup>	4.8 $\pm$ 2.2 <sup>a</sup>	4.1 $\pm$ 2.3 <sup>a</sup>	2.11	0.348
Threonine	Metabolic	–28.2 $\pm$ 3.8	–	–	–	–
Tyrosine	Source	9.2 $\pm$ 2.2 <sup>a</sup>	2.6 $\pm$ 3.0 <sup>b</sup>	2.9 $\pm$ 2.4 <sup>b</sup>	<b>16.4</b>	<b>&lt;0.0001</b>
Valine	Trophic	20.0 $\pm$ 3.2 <sup>a</sup>	21.3 $\pm$ 2.2 <sup>a</sup>	21.7 $\pm$ 2.0 <sup>a</sup>	0.56	0.756
RTP ( $\delta^{15}\text{N}_{\text{Glx}} - \delta^{15}\text{N}_{\text{Phe}}$ )		22.6 $\pm$ 1.8 <sup>a</sup>	22.2 $\pm$ 1.5 <sup>a</sup>	22.8 $\pm$ 1.6 <sup>a</sup>	1.58	0.455
TP <sub>Glx-Phe</sub>		4.4 $\pm$ 0.4 <sup>a</sup>	4.3 $\pm$ 0.3 <sup>a</sup>	4.5 $\pm$ 0.3 <sup>a</sup>	1.32	0.516

in lower and upper beaks. Individual  $\delta^{13}\text{C}_{\text{AA}}$  values ranged widely, from  $-32$  to  $-5$ ‰ for  $\delta^{13}\text{C}_{\text{Leu}}$  and  $\delta^{13}\text{C}_{\text{Gly}}$ , respectively (Fig. 1A). There were no significant isotopic differences amongst the 3 tissues, except the  $\delta^{13}\text{C}_{\text{Thr}}$  value, which was marginally lower in muscle than in beaks (Table 2).

**3.1.3. Amino acid  $\delta^{15}\text{N}$  values.** The method quantified  $\delta^{15}\text{N}_{\text{AA}}$  of 14 AAs in muscle and 11 AAs in beaks.  $\delta^{15}\text{N}_{\text{Hyp}}$  and  $\delta^{15}\text{N}_{\text{Met}}$  cannot be reproducibly measured in muscle, as is the case for  $\delta^{15}\text{N}_{\text{AA}}$  values of Hyp, Ile, Lys, Met and Thr in beaks. In some cases,  $\delta^{15}\text{N}_{\text{AA}}$  values of Pro, Ser, Tyr and Val were difficult to quantify in beaks. Individual  $\delta^{15}\text{N}_{\text{AA}}$  values ranged widely, from  $-28$  to  $23$ ‰ for  $\delta^{15}\text{N}_{\text{Thr}}$  (metabolic AA) and  $\delta^{15}\text{N}_{\text{Ile}}$  (trophic AA), respectively (Table 3). Source AAs (His, Lys, Phe and Tyr) had much lower  $\delta^{15}\text{N}_{\text{AA}}$  values than trophic AAs (Ala, Asx, Glx, Ile, Leu and Val) (Fig. 1B).

Individual  $\delta^{15}\text{N}_{\text{AA}}$  values were consistent amongst tissues, with no differences in 7 AAs, marginally significant differences in 3 AAs and a highly significant difference for Tyr. In the 4 latter cases,  $\delta^{15}\text{N}_{\text{AA}}$  values




Fig. 1. Bulk and individual amino acid (A)  $\delta^{13}\text{C}$  and (B)  $\delta^{15}\text{N}$  values of muscle, lower beak and upper beak from buccal masses of giant warty squids *Kondakovia longimana*. Values are means  $\pm$  SD. \*Significant differences at  $p < 0.05$  (Kruskal-Wallis  $H$ -tests; details in Tables 1 & 2)

were identical in lower and upper beaks but they differed from muscle values. Importantly,  $\delta^{15}\text{N}_{\text{AA}}$  of the 2 canonical AAs—source AA Phe and trophic AA Glx—were not significantly different amongst tissues and, consequently, RTP and  $\text{TP}_{\text{Glx-Phe}}$  were identical when they were calculated using  $\delta^{15}\text{N}_{\text{Glx}}$  and  $\delta^{15}\text{N}_{\text{Phe}}$  values from either muscle, lower or upper beaks (Table 3).

### 3.2. Latitudinal effect on $\delta^{13}\text{C}_{\text{AA}}$ in muscle and beaks

Tissue  $\delta^{13}\text{C}_{\text{Bulk}}$  values grouped giant warty squids in 2 clusters of 5 individuals (muscle:  $-21.3 \pm 0.5$  versus  $-24.2 \pm 0.3\text{‰}$ ; lower beak:  $-20.9 \pm 0.4$  versus  $-24.5 \pm 0.8\text{‰}$ ; upper beak:  $-20.7 \pm 0.4$  versus  $-24.3 \pm 0.6\text{‰}$ ; Mann-Whitney  $U$ -tests, all  $U = 0.0$ , all  $p = 0.009$ ). According to the latitudinal  $\delta^{13}\text{C}_{\text{Bulk}}$  gradient occurring in the southern Indian Ocean (Jaeger et al. 2010), clusters with the higher and lower  $\delta^{13}\text{C}_{\text{Bulk}}$  values corresponded to individuals that grew in subantarctic and Antarctic waters, respectively.

Whatever the tissue and individual AA,  $\delta^{13}\text{C}_{\text{AA}}$  values were always lower in the Antarctic than in the subantarctic group. The difference was significant ( $p < 0.05$ ) in all cases, except for  $\delta^{13}\text{C}_{\text{Thr}}$  in lower beaks ( $U = 5.0$ ,  $p = 0.221$ ) (Fig. 2A). The isotopic difference between each AA of the 2 groups ranged from 1.9 ( $\delta^{13}\text{C}_{\text{Thr}}$ ) to 7.3 $\text{‰}$  ( $\delta^{13}\text{C}_{\text{Gly}}$ ) in muscle, from 1.6 ( $\delta^{13}\text{C}_{\text{Thr}}$ ) to 7.5 $\text{‰}$  ( $\delta^{13}\text{C}_{\text{Gly}}$ ) in lower beak and from 2.9 ( $\delta^{13}\text{C}_{\text{Thr}}$ ) to 8.3 $\text{‰}$  ( $\delta^{13}\text{C}_{\text{Ser}}$ ) in upper beak. AA isotopic differences between Antarctic and subantarctic specimens were consistent across tissues, as illustrated by the positive linear relationship between  $\delta^{13}\text{C}_{\text{AA}}$  differences in 12 AAs from muscle and lower beak (Fig. 2B).

### 3.3. Habitat and trophic position of squids

Darkening or wholly darkened beaks indicated that individuals of the 4 species were either large juvenile or adult squids. Estimated ML ranged from 45 to 239 cm and mean ML values increased in the order giant warty < Dana octopus < giant = colossal squids (Table 1).

**3.3.1. Bulk  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values.** Squids were segregated by both  $\delta^{13}\text{C}_{\text{Bulk}}$  and  $\delta^{15}\text{N}_{\text{Bulk}}$  values of lower beaks. Overall, bulk isotopic values defined 4 distinct isotopic niches (Fig. 3A). Giant warty squids had significantly lower  $\delta^{13}\text{C}_{\text{Bulk}}$  values than Dana octopus, giant and colossal squids. Bulk  $\delta^{15}\text{N}$  values increased in the following order: giant warty = giant < Dana

octopus < colossal squids (Fig. 1A), and, accordingly,  $\text{TP}_{\text{Bulk}}$  followed the same increasing order from 3.9 to 5.9, a 2.0 difference (Table 4).

**3.3.2. AA  $\delta^{13}\text{C}$  values.** Squids were segregated by  $\delta^{13}\text{C}_{\text{AA}}$  of 10 individual AA, with only  $\delta^{13}\text{C}_{\text{Ser}}$  and  $\delta^{13}\text{C}_{\text{Tyr}}$  being not significantly different amongst species. In most cases,  $\delta^{13}\text{C}_{\text{AA}}$  values were lower in the giant warty squid than in other species. Only  $\delta^{13}\text{C}_{\text{Gly}}$  showed species-specific values, ranging from  $-6.4$  (giant warty squid) to  $5.0\text{‰}$  (colossal squid) (Table 4).

**3.3.3. AA  $\delta^{15}\text{N}$  values.** Squids were segregated by  $\delta^{15}\text{N}_{\text{AA}}$  of 10 individual AAs, with only  $\delta^{15}\text{N}_{\text{Val}}$  being




Fig. 2. (A) Bulk and individual amino acid  $\delta^{13}\text{C}$  values of lower beaks from Antarctic and subantarctic giant warty squids *Kondakovia longimana*. Values are means  $\pm$  SD. \*Significant differences at  $p < 0.05$  (Mann-Whitney  $U$ -tests). (B) Linear regression between individual  $\delta^{13}\text{C}_{\text{AA}}$  differences between the Antarctic and subantarctic groups in lower beak and muscle of *K. longimana* ( $y = 0.945x + 0.485$ ,  $r^2 = 0.933$ ,  $F_{1,10} = 139.5$ ,  $p < 0.0001$ ). Black diamond: bulk value; cyan circles: essential AA; red squares: non-essential AAs




Fig. 3. Isotopic niches of oceanic squids using (A)  $\delta^{13}\text{C}_{\text{Bulk}}$  and  $\delta^{15}\text{N}_{\text{Bulk}}$  values and (B) representative  $\delta^{13}\text{C}_{\text{AA}}$  and  $\delta^{15}\text{N}_{\text{AA}}$  values of their lower beaks. Both individual values and species mean values ( $\pm$  SD) are shown. RTP: relative trophic position

not significantly different amongst species. In many cases, individual  $\delta^{15}\text{N}_{\text{AA}}$  values defined 2 groups, with giant and giant warty squids having lower values than colossal and Dana octopus squids.  $\delta^{15}\text{N}_{\text{Glx}}$  values increased in the order giant = giant warty < Dana octopus < colossal squids, and  $\delta^{15}\text{N}_{\text{Phe}}$  values also increased in a similar order: giant = giant warty  $\leq$  Dana octopus < colossal squids (Fig. 4). RTP, and hence  $\text{TP}_{\text{Glx-Phe}}$ , clustered into 2 groups, with giant and giant warty squids having lower values than the colossal and Dana octopus squids. Average  $\text{TP}_{\text{Glx-Phe}}$  ranged from 4.2 to 4.8, thus encompassing a 0.6 difference (Table 4).

**3.3.4. Comparing  $\text{TP}_{\text{Bulk}}$  and  $\text{TP}_{\text{Glx-Phe}}$ .** When pooling all individual squids,  $\text{TP}_{\text{Bulk}}$  estimates did not fit well with  $\text{TP}_{\text{Glx-Phe}}$  (Fig. 5A). At the species level,  $\text{TP}_{\text{Bulk}}$  and  $\text{TP}_{\text{Glx-Phe}}$  values were not different for Dana octopus and giant squids, but they differed signifi-




Fig. 4. (A) Bulk  $\delta^{15}\text{N}$ , (B) RTP and (C)  $\delta^{15}\text{N}_{\text{Phe}}$  values of lower beaks from oceanic squids from Kerguelen waters. Values are means  $\pm$  SD. Kruskal-Wallis  $H$ -tests and Conover-Inman tests for pairwise comparisons were performed to compare values from the 4 squids; different lower-case letters indicate statistical differences at  $p < 0.05$  (details in Table 3). RTP: relative trophic position; *Konda.*: *Kondakovia longimana*; *Archi.*: *Architeuthis dux*; *Tanin.*: *Taningia danae*; *Meson.*: *Mesonychoteuthis hamiltoni*

cantly for colossal and giant warty squids. The  $\text{TP}_{\text{Bulk}}$  estimate of the latter was lower than its  $\text{TP}_{\text{Glx-Phe}}$  value, while  $\text{TP}_{\text{Bulk}}$  of the colossal squid was noticeably higher than its  $\text{TP}_{\text{Glx-Phe}}$  (Table 4). Interestingly,  $\text{TP}_{\text{Bulk}}$  values were more positively related to baseline



Table 4. Bulk and individual amino acid  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of lower beaks, and calculated relative trophic position (RTP) and estimated trophic position (TP) of *Architeuthis dux* (giant squid), *Kondakovia longimana* (giant warty squid), *Mesonychoteuthis hamiltoni* (colossal squid) and *Taningia danae* (Dana octopus squid) from subantarctic Kerguelen waters. Values are means  $\pm$  SD. Kruskal-Wallis  $H$ -tests and Conover-Inman tests for pairwise comparisons were performed to compare  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values from the 4 squids (values not sharing the same superscript letter are statistically different), and paired  $t$ -tests were performed to compare  $\text{TP}_{\text{Glx-Phe}}$  and  $\text{TP}_{\text{Bulk}}$  of each squid species. Significant differences ( $p < 0.05$ ) are highlighted in **bold**. (–) No data

Amino acid	Isotopic values (‰)	<i>Architeuthis dux</i> (n = 6)	<i>Kondakovia longimana</i> (n = 5)	<i>Mesonychoteuthis hamiltoni</i> (n = 10)	<i>Taningia danae</i> (n = 10)	Kruskal-Wallis $H$	$H$ -tests p
Bulk	$\delta^{13}\text{C}$	$-17.9 \pm 0.5^a$	$-21.8 \pm 1.4^b$	$-18.6 \pm 1.6^a$	$-18.6 \pm 1.0^a$	<b>11.7</b>	<b>0.008</b>
	$\delta^{15}\text{N}$	$7.6 \pm 0.5^a$	$6.1 \pm 0.3^a$	$12.9 \pm 1.3^b$	$9.8 \pm 1.6^c$	<b>23.6</b>	<b>&lt;0.0001</b>
Alanine	$\delta^{13}\text{C}$	$-16.5 \pm 1.0^a$	$-23.4 \pm 2.2^b$	$-17.2 \pm 1.4^a$	$-17.2 \pm 2.6^a$	<b>12.6</b>	<b>0.006</b>
	$\delta^{15}\text{N}$	$21.1 \pm 1.1^a$	$19.6 \pm 0.8^a$	$26.2 \pm 0.8^b$	$27.2 \pm 2.1^c$	<b>22.8</b>	<b>&lt;0.0001</b>
Aspartic acid	$\delta^{13}\text{C}$	$-14.4 \pm 0.6^{a,b}$	$-17.2 \pm 1.9^a$	$-13.4 \pm 1.7^b$	$-13.4 \pm 3.1^b$	<b>9.57</b>	<b>0.023</b>
	$\delta^{15}\text{N}$	$17.3 \pm 1.2^a$	$17.4 \pm 0.6^a$	$21.0 \pm 0.8^b$	$20.3 \pm 2.3^b$	<b>16.2</b>	<b>0.001</b>
Glutamic acid	$\delta^{13}\text{C}$	$-16.3 \pm 0.3^a$	$-19.8 \pm 1.8^b$	$-19.1 \pm 1.3^b$	$-18.5 \pm 0.7^b$	<b>15.4</b>	<b>0.002</b>
	$\delta^{15}\text{N}$	$21.2 \pm 1.4^a$	$21.2 \pm 0.9^a$	$26.9 \pm 1.1^b$	$25.4 \pm 2.4^c$	<b>20.3</b>	<b>&lt;0.0001</b>
Glycine	$\delta^{13}\text{C}$	$-0.4 \pm 1.4^a$	$-6.4 \pm 1.3^b$	$5.0 \pm 2.8^c$	$-3.7 \pm 1.7^d$	<b>25.3</b>	<b>&lt;0.0001</b>
	$\delta^{15}\text{N}$	$-3.0 \pm 1.3^a$	$-8.0 \pm 1.9^b$	$9.3 \pm 0.7^c$	$0.6 \pm 1.5^d$	<b>27.3</b>	<b>&lt;0.0001</b>
Histidine	$\delta^{13}\text{C}$	$-7.7 \pm 1.7^a$	$-11.6 \pm 2.8^b$	$-14.8 \pm 3.8^b$	$-6.3 \pm 2.8^a$	<b>20.6</b>	<b>&lt;0.0001</b>
	$\delta^{15}\text{N}$	$4.7 \pm 0.8^{a,b}$	$5.0 \pm 1.2^{a,b}$	$3.4 \pm 1.6^a$	$5.6 \pm 2.0^b$	<b>7.85</b>	<b>0.049</b>
Leucine	$\delta^{13}\text{C}$	$-26.4 \pm 0.6^a$	$-32.5 \pm 1.5^b$	$-28.7 \pm 1.8^c$	$-25.4 \pm 1.6^a$	<b>20.8</b>	<b>&lt;0.0001</b>
	$\delta^{15}\text{N}$	$22.8 \pm 2.4^a$	$22.2 \pm 0.9^a$	$24.6 \pm 1.2^b$	$25.9 \pm 1.8^b$	<b>13.4</b>	<b>0.004</b>
Phenylalanine	$\delta^{13}\text{C}$	$-25.2 \pm 0.9^a$	$-29.4 \pm 1.9^b$	$-28.7 \pm 1.6^b$	$-25.0 \pm 2.0^a$	<b>21.2</b>	<b>&lt;0.0001</b>
	$\delta^{15}\text{N}$	$-0.6 \pm 0.6^a$	$-0.4 \pm 0.6^{a,b}$	$3.0 \pm 1.3^c$	$0.8 \pm 1.6^b$	<b>18.0</b>	<b>&lt;0.0001</b>
Proline	$\delta^{13}\text{C}$	$-17.6 \pm 0.8^a$	$-21.5 \pm 2.0^b$	$-17.1 \pm 1.8^a$	$-18.5 \pm 1.8^a$	<b>11.6</b>	<b>0.009</b>
	$\delta^{15}\text{N}$	$20.5 \pm 2.2^{a,b}$	$20.2 \pm 1.2^a$	$28.0 \pm 2.6^c$	$22.7 \pm 2.4^b$	<b>18.0</b>	<b>&lt;0.0001</b>
Serine	$\delta^{13}\text{C}$	$-3.7 \pm 2.3^a$	$-6.1 \pm 2.6^a$	$-5.3 \pm 3.1^a$	$-4.2 \pm 2.5^a$	3.58	0.310
	$\delta^{15}\text{N}$	$-3.2 \pm 1.7^a$	$4.0 \pm 1.6^b$	$3.9 \pm 2.6^b$	$3.5 \pm 3.0^b$	<b>13.3</b>	<b>0.004</b>
Threonine	$\delta^{13}\text{C}$	$-5.2 \pm 1.4^{a,b}$	$-10.8 \pm 1.6^c$	$-4.1 \pm 3.3^a$	$-7.2 \pm 3.2^b$	<b>12.4</b>	<b>0.006</b>
	$\delta^{15}\text{N}$	–	–	–	–	–	–
Tyrosine	$\delta^{13}\text{C}$	$-23.4 \pm 0.6^a$	$-24.1 \pm 1.7^a$	$-24.6 \pm 1.7^a$	$-22.9 \pm 2.8^a$	2.51	0.473
	$\delta^{15}\text{N}$	$5.0 \pm 1.4^{a,b}$	$3.5 \pm 0.4^a$	$5.3 \pm 1.9^{a,b}$	$6.3 \pm 2.0^b$	<b>7.88</b>	<b>0.048</b>
Valine	$\delta^{13}\text{C}$	$-20.6 \pm 1.0^a$	$-27.0 \pm 2.9^b$	$-16.0 \pm 2.7^c$	$-20.0 \pm 1.8^a$	<b>20.7</b>	<b>&lt;0.0001</b>
	$\delta^{15}\text{N}$	$18.0 \pm 1.9^a$	$19.4 \pm 1.4^a$	$19.0 \pm 3.8^a$	$18.4 \pm 4.1^a$	1.19	0.755
RTP ( $\delta^{15}\text{N}_{\text{Glx}} - \delta^{15}\text{N}_{\text{Phe}}$ )	$\delta^{15}\text{N}$	$21.8 \pm 1.5^a$	$21.7 \pm 0.8^a$	$23.9 \pm 1.5^b$	$24.6 \pm 1.4^b$	<b>15.2</b>	<b>0.002</b>
$\text{TP}_{\text{Glx-Phe}}$		$4.3 \pm 0.3^a$	$4.2 \pm 0.2^a$	$4.7 \pm 0.3^b$	$4.8 \pm 0.3^b$	<b>15.2</b>	<b>0.002</b>
$\text{TP}_{\text{Bulk}}$		$4.3 \pm 0.5^a$	$3.9 \pm 0.1^a$	$5.9 \pm 0.4^b$	$5.0 \pm 0.5^c$	<b>23.6</b>	<b>&lt;0.0001</b>
Paired $t$ -tests	$t$	0.49	<b>-8.69</b>	<b>9.44</b>	1.24		
	p	0.643	<b>0.001</b>	<b>&lt;0.0001</b>	0.246		
Difference ( $\text{TP}_{\text{Bulk}} - \text{TP}_{\text{Glx-Phe}}$ )		$0.1 \pm 0.5^a$	$-0.3 \pm 0.1^b$	$1.3 \pm 0.4^c$	$0.2 \pm 0.5^a$	<b>20.9</b>	<b>&lt;0.0001</b>

$\delta^{15}\text{N}_{\text{Phe}}$  values ( $y = 0.36x + 4.63$ ,  $r^2 = 0.615$ ,  $F_{1,29} = 46.4$ ,  $p < 0.0001$ ) (Fig. 5B) than to trophic  $\delta^{15}\text{N}_{\text{CSIA}}$  values (RTP;  $y = 0.27x - 1.20$ ,  $r^2 = 0.310$ ,  $F_{1,29} = 13.0$ ,  $p = 0.001$ ).

$\text{TP}_{\text{Bulk}}$  and  $\text{TP}_{\text{Glx-Phe}}$  were not correlated with ML within each squid species (data not shown), thus suggesting no ontogenetic dietary shift within the investigated size ranges. Estimated ML of giant warty squids collected in Kerguelen waters in 2014–2015 ( $n = 7$ , first data set) and 1997–2001 ( $n = 5$ , second data set) were identical, as were their  $\delta^{15}\text{N}_{\text{Bulk}}$ ,  $\text{TP}_{\text{Bulk}}$ ,  $\delta^{15}\text{N}_{\text{Phe}}$ ,  $\delta^{15}\text{N}_{\text{Glx}}$ , RTP and  $\text{TP}_{\text{Glx-Phe}}$  values (Mann-Whitney  $U$ -tests, all  $p \geq 0.329$ ).

#### 4. DISCUSSION

This study presents an innovative method to investigate the trophic ecology of cephalopods that complements the use of bulk isotopic values of their beaks (Cherel & Hobson 2005). It is the first to validate and test CSIA-AA on cephalopods by measuring their beak  $\delta^{13}\text{C}_{\text{AA}}$  and  $\delta^{15}\text{N}_{\text{AA}}$  values. Since the method focuses on AA from proteins, it eliminates problems due to the presence of other organic compounds that may affect bulk isotopic values. For  $\delta^{15}\text{N}$ , CSIA-AA discards chitin during the analytical procedure, thus removing the chitin effect that lowers




Fig. 5. Relationships between (A) TP<sub>Bulk</sub> and TP<sub>Glx-Phe</sub>, and (B) TP<sub>Bulk</sub> and  $\delta^{15}\text{N}_{\text{Phe}}$  values of lower beaks from 4 oceanic squids from Kerguelen waters. Values are means  $\pm$  SD. TP: trophic position

beak  $\delta^{15}\text{N}_{\text{Bulk}}$  values (Cherel et al. 2009a). Additionally, the most important applications of  $\delta^{15}\text{N}$  CSIA-AA to date is to disentangle the  $\delta^{15}\text{N}$  baseline effect from the  $\delta^{15}\text{N}$  trophic effect by simultaneously measuring  $\delta^{15}\text{N}_{\text{Source AA}}$  and  $\delta^{15}\text{N}_{\text{Trophic AA}}$  in the consumer itself, thus allowing better TP<sub>CSIA</sub> estimates than using  $\delta^{15}\text{N}_{\text{Bulk}}$  values (McMahon & McCarthy 2016).

#### 4.1. Validating the use of $\delta^{13}\text{C}_{\text{AA}}$ and $\delta^{15}\text{N}_{\text{AA}}$ in cephalopod beaks

As expected,  $\delta^{13}\text{C}_{\text{AA}}$  and  $\delta^{15}\text{N}_{\text{AA}}$  values were measured on 14–15 standard AA isolated from muscle of the giant warty squid, thus validating the analytical procedure on cephalopod tissues. Smaller numbers

of AAs were measured in beaks, which can be explained by tissue-specific AA composition (Hunt & Nixon 1981, Miserez et al. 2007). Data from giant warty squids may be generalized to other cephalopods, because beaks of Dana octopus, and colossal and giant squids gave similar results, which were therefore consistent across species and across beaks with different sclerotized levels that reflect different amounts of chitin relative to proteins (Rubin et al. 2010). To sum up, beak  $\delta^{13}\text{C}_{\text{AA}}$  can be confidently measured on 12 AAs that include 5 essential (His, Leu, Phe, Thr, Val) and 7 non-essential (Ala, Asx, Glx, Gly, Pro, Ser, Tyr) AAs, and beak  $\delta^{15}\text{N}_{\text{AA}}$  can be quantified on at least 7 AAs that include 2 source (His, Phe) and 4 trophic (Ala, Asx, Glx, Leu) AAs.

Another major finding was the similarity of  $\delta^{13}\text{C}_{\text{AA}}$  and  $\delta^{15}\text{N}_{\text{AA}}$  values amongst tissues. Lower and upper beaks showed no AA isotopic differences, and beaks and muscle had either identical or slightly different AA isotopic values. The only exception is  $\delta^{15}\text{N}_{\text{Tyr}}$ , which was much higher in muscle, and we have no explanation for that difference. Three consequences of the inter-tissue isotopic comparison are notable. (1) Since CSIA-AA overcomes the chitin effect on beaks, the method may be successfully applied to the gladius (Ruiz-Cooley et al. 2013), another chitin-containing hard structure (Hunt & Nixon 1981), whose morphology allows serial sampling to isotopically reconstruct the past individual trophic history of squids (Cherel et al. 2009a, Ruiz-Cooley et al. 2010, Lorrain et al. 2011). (2) Beak (hard tissue) is as representative as muscle (soft tissue) to investigate the isotopic ecology of cephalopods. This paves the way to use the numerous accumulated beaks sorted from predator stomachs to gather useful biological information on oceanic cephalopods. (3) Since  $\delta^{15}\text{N}_{\text{Glx}}$  and  $\delta^{15}\text{N}_{\text{Phe}}$  did not vary amongst tissues with a similar time integration period, RTP and TP<sub>Glx-Phe</sub> were not tissue-specific and accurate TP<sub>Glx-Phe</sub> estimates can be calculated either from beak or muscle values.

#### 4.2. Testing the usefulness of beak $\delta^{13}\text{C}_{\text{AA}}$ and $\delta^{15}\text{N}_{\text{AA}}$ of cephalopods

**4.2.1. Foraging habitat and beak  $\delta^{13}\text{C}_{\text{AA}}$ .** The most important application of  $\delta^{13}\text{C}_{\text{Bulk}}$  of marine consumers is to define their isotopic habitats as a proxy of their foraging habitats. Tissue  $\delta^{13}\text{C}_{\text{Bulk}}$  clustered giant warty squids into an Antarctic and a subantarctic group, thus allowing us to test the ability of  $\delta^{13}\text{C}_{\text{AA}}$  to discriminate the squid feeding grounds. All  $\delta^{13}\text{C}_{\text{AA}}$

values from the 3 tissues were lower in the Antarctic than subantarctic group. This confirms—for muscle and beaks of cephalopods—the results obtained on blood and feathers of penguins at different spatial oceanographic scales (Lorrain et al. 2009, Polito et al. 2017), and emphasizes the usefulness of  $\delta^{13}\text{C}_{\text{AA}}$  to depict foraging latitudes and migration patterns of consumers from the Southern Ocean. Thus,  $\delta^{13}\text{C}_{\text{AA}}$  of accumulated beaks can also help with reconstructing the foraging habitats of cephalopod eaters (e.g. albatrosses, sharks) over the weeks/months preceding sampling.

Not all AAs were equally efficient to differentiate feeding grounds. Surprisingly, essential AAs discriminated less well the 2 latitudinal habitats than non-essential AAs, with Gly being the most discriminant AAs. Leu was the single exception, because it grouped with the non-essential AAs (Fig. 2B). The data compare well with a previous investigation on penguins showing that  $\delta^{13}\text{C}_{\text{AA}}$  of all measured AAs vary with  $\delta^{13}\text{C}_{\text{Bulk}}$  and latitudes (Lorrain et al. 2009). Some previous studies focused on essential AAs (Polito et al. 2017), the rationale being that essential AAs tracked  $\delta^{13}\text{C}$  baseline levels, because many non-essential AAs are trophically  $^{13}\text{C}$  enriched (McMahon et al. 2013, 2015). Here, there was no differential trophic effect between the 2 groups of giant warty squids, since RTP and  $\text{TP}_{\text{Glx-Phe}}$  of Antarctic and subantarctic specimens were not significantly different, whatever the tissue considered (Mann-Whitney  $U$ -tests, all  $p \geq 0.175$ ). Consequently, the usefulness of the non-essential Gly was tested in the second set of samples. Amongst the 4 squid species, lower beak  $\delta^{13}\text{C}_{\text{Gly}}$  was positively and linearly related to  $\delta^{15}\text{N}_{\text{Phe}}$  ( $y = 1.63x - 2.38$ ,  $r^2 = 0.417$ ,  $F_{1,29} = 20.7$ ,  $p < 0.0001$ ), but it did not correlate with RTP, meaning that  $\delta^{13}\text{C}_{\text{Gly}}$  was more linked to isotopic baseline than to trophic enrichment. Consequently, our data highlight that  $\delta^{13}\text{C}_{\text{AA}}$  of non-essential AAs can be helpful to delineate foraging habitats of consumers, and this merits further investigations on other animal models living in different ecosystems.

**4.2.2. Estimating  $\text{TP}_{\text{CSIA}}$  using beak  $\delta^{15}\text{N}_{\text{AA}}$ .** RTP and  $\text{TP}_{\text{Glx-Phe}}$  estimates grouped the 4 oceanic squids into 2 clusters, with giant and giant warty squids having lower values than colossal and Dana octopus squids. This contrasts with  $\delta^{15}\text{N}_{\text{Bulk}}$  values depicting 3 different  $\text{TP}_{\text{Bulk}}$  estimates. Comparing RTP and  $\text{TP}_{\text{Glx-Phe}}$  to  $\delta^{15}\text{N}_{\text{Bulk}}$  and  $\text{TP}_{\text{Bulk}}$  emphasizes the ability of CSIA-AA to quantify and disentangle isotopic  $\delta^{15}\text{N}$  baseline from trophic  $^{15}\text{N}$  enrichment (McMahon et al. 2013, Ohkouchi et al. 2017). This is well illustrated by the positive correlation between squid  $\text{TP}_{\text{Bulk}}$  and

$\delta^{15}\text{N}_{\text{Phe}}$  that indicates a consistent baseline effect on  $\delta^{15}\text{N}_{\text{Bulk}}$  and  $\text{TP}_{\text{Bulk}}$  values.  $\text{TP}_{\text{Glx-Phe}}$  and  $\text{TP}_{\text{Bulk}}$  were identical for Dana octopus and giant squids, slightly different for giant warty squid, but strongly differed for colossal squid.  $\text{TP}_{\text{Bulk}}$  of the latter species was previously estimated at 6.1 (Cherel et al. 2008), a value close to 5.9 reported here. By contrast, its  $\text{TP}_{\text{Glx-Phe}}$  was 4.7, thus lowering TP by  $\sim 1.3$ , which is a highly relevant difference to assess the role of colossal squid within the oceanic ecosystem both as a predator and prey. Overestimation of  $\delta^{15}\text{N}_{\text{Bulk}}$  and  $\text{TP}_{\text{Bulk}}$  due to a baseline effect is also the likely explanation of some high values recorded in the literature for deep-sea oceanic squids (e.g. gonatids, histioteuthids), which include small and delicate forms (e.g. chiroteuthids, mastigoteuthids) (Cherel & Hobson 2005, Cherel et al. 2008, 2009b, Guerreiro et al. 2015, Golikov et al. 2018). Clearly, the issue merits further investigations using CSIA-AA to compare thoroughly  $\text{TP}_{\text{Bulk}}$  with  $\text{TP}_{\text{CSIA}}$  estimates.

Knowledge on the food of colossal, Dana octopus, giant and giant warty squids remains very limited. The few dietary information collected opportunistically from cephalopods eaten by sperm whales showed that both colossal and giant warty squids prey primarily upon mesopelagic fish (Lubimova 1985). Accordingly,  $\text{TP}_{\text{Bulk}}$  estimates of most Kerguelen myctophids, the main mesopelagic fish biomass of the Southern Ocean, are lower (3.3–3.9) than squid  $\text{TP}_{\text{Glx-Phe}}$  (4.2–4.8), and the latter compares well with  $\text{TP}_{\text{Bulk}}$  estimates (4.3–4.8) of predators that are known to feed primarily on myctophids (blue petrel, king penguin, southern elephant seal and Antarctic fur seal) (Cherel et al. 2008, 2010, 2017). Hence, the largest invertebrates living on Earth are not apex predators, but, instead, they exploit mesopelagic fish that constitutes the highest oceanic micronektonic biomass available in the Southern Ocean and worldwide (Kozlov 1995, Irigoien et al. 2014).

**4.2.3. Comparing the isotopic niche using bulk and AA  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ .** In Kerguelen waters, beak  $\delta^{13}\text{C}_{\text{Bulk}}$  and  $\delta^{15}\text{N}_{\text{Bulk}}$  values depict species-specific isotopic niches, with each sympatric squid using a unique combination amongst 2 isotopic habitats and 3  $\text{TP}_{\text{Bulk}}$  (Fig. 3A). CSIA-AA challenged this traditional approach. RTP and  $\text{TP}_{\text{Glx-Phe}}$  specified that trophic segregation operates at 2 different TP, and  $\delta^{15}\text{N}_{\text{Phe}}$  depicted different baseline levels corresponding to 3 foraging habitats. The use of Gly (the most discriminant AA for habitat; see above) further detailed spatial segregation with 4 different  $\delta^{13}\text{C}_{\text{Gly}}$  values. Consequently, a contrasted and more precise picture emerged when using CSIA-AA. Again, iso-

topic niches were species-specific, but along 4 isotopic habitats and 2 TP<sub>Gly-Phe</sub> (Fig. 3B).

Three consequences of the comparison between bulk and AA isotopic values of squid beaks are notable. (1) A principle in isotopic ecology is that isotopic differences carry relevant biological information, but a lack of isotopic difference does not always correspond to identical ecological features. This latter issue is exemplified by the squid foraging habitats. While  $\delta^{13}\text{C}_{\text{Bulk}}$  values of colossal, Dana octopus and giant squids suggested identical foraging grounds, both  $\delta^{13}\text{C}_{\text{Gly}}$  and  $\delta^{15}\text{N}_{\text{Phe}}$  depicted 3 contrasting feeding habitats, thus underlining how difficult the biological interpretation of isotopic data can be. (2) Within that context, it is notable that most animal isotopic studies are based on bulk analysis, a few on  $\delta^{15}\text{N}$  CSIA-AA, a very few on  $\delta^{13}\text{C}$  CSIA-AA and almost none on both  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  CSIA-AA (but see Petzke et al. 2005, Jarman et al. 2017, Pomerleau et al. 2017). The present work underlines that the isotopic method is at its best when including concomitant measurements of bulk and AA  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  analyses on the same samples. (3) A recurrent limitation of the bulk isotopic method is the availability (or not) of marine isoscapes to help interpret isotopic data in terms of meaningful biological information (Graham et al. 2010, McMahon et al. 2013). The problem is even more difficult when using CSIA-AA, due to the complete lack of information. We recommend that future studies aim at constructing maps of the geographical distribution of  $\delta^{13}\text{C}_{\text{AA}}$  values of essential and non-essential AAs, and for  $\delta^{15}\text{N}_{\text{AA}}$  values of source AAs at spatial scales that are ecologically relevant to the studied animals.

### 4.3. Perspectives

The presence of chitin is not restricted to cephalopod beaks. It is one of the most abundant macromolecules in the biosphere, being a main component of arthropods, the most diverse and successful animals on Earth. Arthropod exoskeleton is made of cuticle, which consists of varying amounts of protein and chitin, with the latter representing up to 40% of its dry mass (Merzendorfer & Zimoch 2003). Many isotopic investigations include arthropods, and, owing to their small size, isotopic measurements were generally made on whole organisms that include exoskeleton. Arthropod isotopic studies overlook the deleterious effect of chitin, which applies to both crustaceans and insects (Søreide & Nygard 2012, Perkins et al. 2013). Consequently, arthropod

TP<sub>Bulk</sub> values are likely to be systematically underestimated (e.g. Chikaraishi et al. 2011, Steffan et al. 2013), which can alter the description and functioning of trophic relationships. Calculation of consumer TP<sub>Bulk</sub> requires the use of  $\delta^{15}\text{N}_{\text{Bulk}}$  value of a food web baseline that is, in many cases, a primary consumer. Herbivorous copepods and euphausiids are the mostly commonly used marine organisms (e.g. Marsh et al. 2017, McClain-Counts et al. 2017), thus propagating the isotopic error associated with low  $\delta^{15}\text{N}_{\text{Bulk}}$  values to consumer TP<sub>Bulk</sub> estimates through the food web. Finally, some maps of marine  $\delta^{15}\text{N}_{\text{Bulk}}$  isoscapes are modeled using zooplankton data sets (Graham et al. 2010, McMahon et al. 2013), with the direct potential biological outcome of erroneous interpretation of animal movements.

To conclude, CSIA-AA on sclerotized beaks is a powerful tool to investigate trophic interactions of cephalopods. It has a great potential due to the high number of beaks that accumulate in predator stomachs. CSIA-AA is intrinsically effective to bypass the deleterious effect of non-proteinaceous compounds (e.g. chitin on  $\delta^{15}\text{N}_{\text{Bulk}}$ , lipids on  $\delta^{13}\text{C}_{\text{Bulk}}$ ), which hamper the biological interpretation of bulk isotopic values. Due to the high chitin content of exoskeleton, CSIA-AA merits further consideration in studies focusing on the ecological role of arthropods in both marine and terrestrial ecosystems.

*Acknowledgements.* The authors thank G. Guillou from the Plateforme Analyses Isotopiques (LIENSs) for stable isotope analysis. They acknowledge the fishery team from the MNHN (Paris) for providing data, fishery observers from the TAAF for collecting samples at sea, the crews and ship-owners of longliners and trawlers operating within the French EEZ around Kerguelen and Crozet Islands, and the French ministry of Agriculture and Food (Direction des Pêches Maritimes et de l'Aquaculture, DPMA) for its financial support. The authors are grateful to the CPER (Contrat de Projet État-Région) and the FEDER (European Regional Development Fund) for funding the IRMS. The IUF (Institut Universitaire de France) is acknowledged for its support to P.B. as a Senior Member.

### LITERATURE CITED

- ✦ Adams NJ, Klages NT (1987) Seasonal variation in the diet of the king penguin (*Aptenodytes patagonicus*) at sub-Antarctic Marion Island. *J Zool* 212:303–324
- ✦ Cherel Y, Duhamel G (2004) Antarctic jaws: cephalopod prey of sharks in Kerguelen waters. *Deep Sea Res I* 51: 17–31
- ✦ Cherel Y, Hobson KA (2005) Stable isotopes, beaks and predators: a new tool to study the trophic ecology of cephalopods, including giant and colossal squids. *Proc R Soc Biol Sci* 272:1601–1607

- Cherel Y, Duhamel G, Gasco N (2004) Cephalopod fauna of subantarctic islands: new information from predators. *Mar Ecol Prog Ser* 266:143–156
- Cherel Y, Ducatez S, Fontaine C, Richard P, Guinet C (2008) Stable isotopes reveal the trophic position and mesopelagic fish diet of female southern elephant seals breeding on the Kerguelen Islands. *Mar Ecol Prog Ser* 370: 239–247
- Cherel Y, Fontaine C, Jackson GD, Jackson CH, Richard P (2009a) Tissue, ontogenic and sex-related differences in  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of the oceanic squid *Todarodes filippovae* (Cephalopoda: Ommastrephidae). *Mar Biol* 156:699–708
- Cherel Y, Ridoux V, Spitz J, Richard P (2009b) Stable isotopes document the trophic structure of a deep-sea cephalopod assemblage including giant octopus and giant squid. *Biol Lett* 5:364–367
- Cherel Y, Fontaine C, Richard P, Labat JP (2010) Isotopic niches and trophic levels of myctophid fishes and their predators in the Southern Ocean. *Limnol Oceanogr* 55: 324–332
- Cherel Y, Xavier JC, de Grissac S, Trouvé C, Weimerskirch H (2017) Feeding ecology, isotopic niche, and ingestion of fishery-related items of the wandering albatross *Diomedea exulans* at Kerguelen and Crozet Islands. *Mar Ecol Prog Ser* 565:197–215
- Chikaraishi Y, Ogawa NO, Ohkouchi N (2010) Further evaluation of the trophic level estimation based on nitrogen isotopic composition of amino acids. In: Ohkouchi N, Tayasu I, Koba K (eds) *Earth, life and isotopes*. Kyoto University Press, Kyoto, p 37–51
- Chikaraishi Y, Ogawa NO, Doi H, Ohkouchi N (2011)  $^{15}\text{N}/^{14}\text{N}$  ratios of amino acids as a tool for studying terrestrial food webs: a case study of terrestrial insects (bees, wasps, and hornets). *Ecol Res* 26:835–844
- Chikaraishi Y, Steffan SA, Ogawa NO, Ishikawa NF, Sasaki Y, Tsuchiya M, Ohkouchi N (2014) High-resolution food webs based on nitrogen isotopic composition of amino acids. *Ecol Evol* 4:2423–2449
- Choy CA, Popp BN, Hannides CCS, Drazen JC (2015) Trophic structure and food resources of epipelagic and mesopelagic fishes in the North Pacific Subtropical Gyre ecosystem inferred from nitrogen isotopic compositions. *Limnol Oceanogr* 60:1156–1171
- Clarke MR (1980) Cephalopoda in the diet of sperm whales of the Southern Hemisphere and their bearing on sperm whale biology. *Discov Rep* 37:1–324
- Clarke MR (1986) *A handbook for the identification of cephalopod beaks*. Clarendon Press, Oxford
- Clarke MR (1996) The role of cephalopods in the world's oceans. *Philos Trans R Soc Lond B Biol Sci* 351:977–1112
- Coll M, Navarro J, Olson RJ, Christensen V (2013) Assessing the trophic position and ecological role of squids in marine ecosystems by means of food-web models. *Deep Sea Res II* 95:21–36
- Docherty G, Jones V, Evershed RP (2001) Practical and theoretical considerations in the gas chromatography/combustion/isotope ratio mass spectrometry  $\delta^{13}\text{C}$  analysis of small polyfunctional compounds. *Rapid Commun Mass Spectrom* 15:730–738
- Fountoulakis M, Lahm HW (1998) Hydrolysis and amino acid composition analysis of proteins. *J Chromatogr A* 826:109–134
- Golikov AV, Ceia FR, Sabirov RM, Zaripova ZI, Blicher ME, Zakharov DV, Xavier JC (2018) Ontogenetic changes in stable isotope ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) values in squid *Gonatus fabricii* (Cephalopoda) reveal its important ecological role in the Arctic. *Mar Ecol Prog Ser* 606:65–78
- Graham BS, Koch PL, Newsome SD, McMahon KW, Aurioules D (2010) Using isoscapes to trace the movements and foraging behaviour of top predators in oceanic ecosystems. In: West JB, Bowen GJ, Dawson TE, Tu KP (eds) *Isoscapes: understanding movement, pattern, and process on Earth through isotope mapping*. Springer, New York, NY, p 299–318
- Guerreiro M, Phillips RA, Cherel Y, Ceia FR, Alvito P, Rosa R, Xavier JC (2015) Habitat and trophic ecology of Southern Ocean cephalopods from stable isotope analyses. *Mar Ecol Prog Ser* 530:119–134
- Hetherington ED, Olson RJ, Drazen JC, Lennert-Cody CE, Balance LT, Kaufmann RS, Popp BN (2017) Spatial food-web structure in the eastern tropical Pacific Ocean based on compound-specific nitrogen isotope analysis of amino acids. *Limnol Oceanogr* 62:541–560
- Hobson KA, Cherel Y (2006) Isotopic reconstruction of marine food webs using cephalopod beaks: new insight from captive raised *Sepia officinalis*. *Can J Zool* 84:766–770
- Hunt S, Nixon M (1981) A comparative study of protein composition in the chitin-protein complexes of the beak, pen, sucker disc, radula and oesophageal cuticle of cephalopods. *Comp Biochem Physiol B* 68:535–546
- Irigoin X, Klevjer TA, Rostad A, Martinez U and others (2014) Large mesopelagic fishes biomass and trophic efficiency in the open ocean. *Nat Commun* 5:3271
- Jaeger A, Lecomte VJ, Weimerskirch H, Richard P, Cherel Y (2010) Seabird satellite tracking validates the use of latitudinal isoscapes to depict predators' foraging areas in the Southern Ocean. *Rapid Commun Mass Spectrom* 24: 3456–3460
- Jarman CL, Larsen T, Hunt T, Lipo C and others (2017) Diet of prehistoric population of Rapa Nui (Easter Island, Chile) shows environmental adaptation and resilience. *Am J Phys Anthropol* 164:343–361
- Kelly JF (2000) Stable isotopes of carbon and nitrogen in the study of avian and mammalian trophic ecology. *Can J Zool* 78:1–27
- Kozlov AN (1995) A review of the trophic role of mesopelagic fish of the family Myctophidae in the Southern Ocean ecosystem. *CCAMLR Sci* 2:71–77
- Kruse S, Pakhomov EA, Hunt BPV, Chikaraishi Y, Ogawa NO, Bathmann U (2015) Uncovering the trophic relationship between *Themisto gaudichaudii* and *Salpa Thompsoni* in the Antarctic Polar Frontal Zone. *Mar Ecol Prog Ser* 529:63–74
- Lehninger AL (1982) *Principles of biochemistry*. Worth Publishers, New York, NY
- Lorrain A, Graham B, Ménard F, Popp B, Bouillon S, van Breugel P, Cherel Y (2009) Nitrogen and carbon isotope values of individual amino acids: a tool to study foraging ecology of penguins in the Southern Ocean. *Mar Ecol Prog Ser* 391:293–306
- Lorrain A, Arguelles J, Alegre A, Bertrand A, Munaron JM, Richard P, Cherel Y (2011) Sequential isotopic signature along gladius highlights contrasted individual foraging strategies of jumbo squid (*Dosidicus gigas*). *PLOS ONE* 6:e22194
- Lubimova TG (1985) Results of Soviet investigation of the distribution and ecology of pelagic squids (Oegopsida) in the Southern Ocean. *Selected Papers SC-CCAMLR* 1985:79–111

- Madigan DJ, Chiang WC, Wallsgrrove NJ, Popp BN and others (2016) Intrinsic tracers reveal recent foraging ecology of giant Pacific bluefin tuna at their primary spawning grounds. *Mar Ecol Prog Ser* 553:253–266
- Marsh JM, Mueter FJ, Iken K, Danielson S (2017) Ontogenetic, spatial and temporal variation in trophic level and diet of Chukchi Sea fishes. *Deep-Sea Res II* 135: 78–94
- McClain-Counts JP, Demopoulos AWJ, Ross SW (2017) Trophic structure of mesopelagic fishes in the Gulf of Mexico revealed by gut content and stable isotope analyses. *Mar Ecol* 38:e12449
- McMahon KW, McCarthy MD (2016) Embracing variability in amino acid  $\delta^{15}\text{N}$  fractionation: mechanisms, implications, and applications for trophic ecology. *Ecosphere* 7: e01511
- McMahon KW, Hamady LL, Thorrold SR (2013) Ocean geochemistry: a review. *Oceanogr Mar Biol Annu Rev* 51:327–374
- McMahon KW, Polito MJ, Abel S, McCarthy MD, Thorrold SR (2015) Carbon and nitrogen isotope fractionation of amino acids in an avian marine predator, the gentoo penguin (*Pygoscelis papua*). *Ecol Evol* 5:1278–1290
- Merzendorfer H, Zimoch L (2003) Chitin metabolism in insects: structure, function and regulation of chitin synthases and chitinases. *J Exp Biol* 206:4393–4412
- Miserez A, Li Y, Waite JH, Zok F (2007) Jumbo squid beaks: inspiration for design of robust organic composites. *Acta Biomater* 3:139–149
- Navarro J, Coll M, Somes CJ, Olson RJ (2013) Trophic niche of squids: insights from isotopic data in marine systems worldwide. *Deep Sea Res II* 95:93–102
- Nielsen JM, Popp BN, Winder M (2015) Meta-analysis of amino acid stable nitrogen isotope ratios for estimating trophic position in marine organisms. *Oecologia* 178: 631–642
- Ohkouchi N, Chikaraishi Y, Close HG, Fry B and others (2017) Advances in the application of amino acid nitrogen isotopic analysis in ecological and biogeochemical studies. *Org Geochem* 113:150–174
- Pakhomov EA, Henschke N, Hunt BPV, Stowasser G, Cherel Y (2019) Utility of salps as a baseline proxy for food web studies. *J Plankton Res* 41:3–11
- Perkins MJ, McDonald RA, van Veen FJF, Kelly SD, Rees G, Bearhop S (2013) Important impacts of tissue selection and lipid extraction on ecological parameters derived from stable isotope ratios. *Methods Ecol Evol* 4:944–953
- Petzke KJ, Boeing H, Klaus S, Metges CC (2005) Carbon and nitrogen stable isotopic composition of hair protein and amino acids can be used as biomarkers for animal-derived dietary protein intake in humans. *J Nutr* 135: 1515–1520
- Polito MJ, Hinke JT, Hart T, Santos M, Houghton LA, Thorrold SR (2017) Stable isotope analyses of feather amino acids identify penguin migration at ocean basin scales. *Biol Lett* 13:20170241
- Pomerleau C, Heide-Jørgensen MP, Ferguson SH, Stern HL, Høyer JL, Stern GA (2017) Reconstruction variability in West Greenland ocean biogeochemistry and bowhead whale (*Balaena mysticetus*) food web structure using amino acid isotope ratios. *Polar Biol* 40:2225–2238
- Post DM (2002) Using stable isotopes to estimate trophic position: models, methods, and assumptions. *Ecology* 83: 703–718
- Queiros JP, Cherel Y, Ceia FR, Hilario A, Roberts J, Xavier JC (2018) Ontogenetic changes in habitat and trophic ecology in the Antarctic squid *Kondakovia longimana* derived from isotopic analysis on beaks. *Polar Biol* 41: 2409–2421
- Rodhouse PG (1990) Cephalopod fauna of the Scotia Sea at South Georgia: potential for commercial exploitation and possible consequences. In: Kerry KR, Hempel G (eds) *Antarctic ecosystems: ecological change and conservation*. Springer Verlag, Berlin, p 289–298
- Roelvelde MAC (2000) Giant squid beaks: implications for systematics. *J Mar Biol Assoc UK* 80:185–187
- Rubin DJ, Miserez A, Waite JH (2010) Diverse strategies of protein sclerotization in marine invertebrates: structure-property relationships in natural biomaterials. *Adv Insect Physiol* 38:75–133
- Ruiz-Cooley RI, Markaida U, Gendron D, Aguinaga S (2006) Stable isotopes in jumbo squid (*Dosidicus gigas*) beaks to estimate its trophic position: comparison between stomach contents and stable isotopes. *J Mar Biol Assoc UK* 86: 437–445
- Ruiz-Cooley RI, Villa EC, Gould WR (2010) Ontogenetic variation of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  recorded in the gladius of the jumbo squid *Dosidicus gigas*: geographic differences. *Mar Ecol Prog Ser* 399:187–198
- Ruiz-Cooley RI, Ballance LT, McCarthy MD (2013) Range expansion of the jumbo squid in the NE Pacific:  $\delta^{15}\text{N}$  decrypts multiple origins, migration and habitat use. *PLOS ONE* 8:e59651
- Schimmelmann A (2011) Carbon, nitrogen and oxygen stable isotope ratios in chitin. In: Gupta NS (ed) *Chitin. Topics in geobiology*, Vol 34. Springer, Dordrecht, p 81–103
- Søreide JE, Nygard H (2012) Challenges using stable isotopes for estimating trophic levels in marine amphipods. *Polar Biol* 35:447–453
- Steffan SA, Chikaraishi Y, Horton DR, Ohkouchi N and others (2013) Trophic hierarchies illuminated via amino acid isotopic analysis. *PLOS ONE* 8:e76152
- Styring AK, Kuhl A, Knowles TDJ, Fraser RA, Bogaard A, Evershed RP (2012) Practical considerations in the determination of compound-specific amino acid  $\delta^{15}\text{N}$  values in animal and plant tissues by gas chromatography-combustion-isotope ratio mass spectrometry, following derivatization to their N-acetylisopropyl esters. *Rapid Commun Mass Spectrom* 26:2328–2334
- Vanderklift A, Ponsard S (2003) Sources of variation in consumer-diet  $\delta^{15}\text{N}$  enrichments: a meta-analysis. *Oecologia* 136:169–182
- Walsh RG, He S, Yarnes CT (2014) Compound-specific  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  analysis of amino acids: a rapid, chloroformate-based method for ecological studies. *Rapid Commun Mass Spectrom* 28:96–108
- Webb S, Hedges REM, Simpson SJ (1998) Diet quality influences the  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  of locusts and their biochemical components. *J Exp Biol* 201:2903–2911
- Xavier JC, Cherel Y (2009) Cephalopod beak guide for the Southern Ocean. British Antarctic Survey, Cambridge
- Xavier JC, Allcock AL, Cherel Y, Lipinski MR and others (2015) Future challenges in cephalopod research. *J Mar Biol Assoc UK* 95:999–1015